

GOVERNMENT SUBSIDIZED INOCULATION

For residents of Tokyo's 23 wards (April, 2016)

vaccine	aid recipients	standard inoculation age and doses	the age coupon delivered
Hib hemophilus influenza type b vaccine	initial: age of 2mo-up to 5yr ----- booster: age of 2mo-up to 5yr	3 times: age of 2mo-6mo (4-8wk interval each) ----- once: 7-13mo after completing initial inoculation	at the age of 2mo (3+1 coupons)
PCV 13 pneumococcal polysaccharide conjugated vaccine	initial: age of 2mo-up to 5yr ----- booster: age of 2mo-up to 5yr	3 times: age of 2mo-6mo (27day interval each) ----- once: age of 1yr/ 60 days after completing initial inoculation	at the age of 2mo (3+1 coupons)
BCG (UPPER OUTER ARM ONLY WITH COUPON) bacille de calmette guerin (against tuberculosis)	up to age of 1yr	once: age of 5mo-up to 8mo	at the age of 3mo (1 coupon)
DPT-IPV: primary period (quadrivalent) diphtheria, tetanus and acellular pertussis-IPV	initial: age of 3mo-up to 90mo ----- booster: age of 3mo-up to 90mo	3 times: age of 3mo-12mo (3-8wk interval each) ----- once: 12-18mo after completing initial inoculation	at the age of 3mo (3 coupons) ----- at the age of 18mo (1 coupon)
DPT: primary period (trivalent) diphtheria, tetanus and acellular pertussis	initial: age of 3mo-up to 90mo ----- booster: age of 3mo-up to 90mo	3 times: age of 3mo-12mo (3-8wk interval each) ----- once: 12-18mo after completing initial inoculation	inform Health Center if desired
IPV inactivated polio vaccine	initial: age of 3mo-up to 90mo ----- booster: age of 3mo-up to 90mo	3 times: age of 3mo-12mo (3-8wk interval each) ----- once: 12-18mo after completing initial inoculation	inform Health Center if desired
DT: secondary period diphtheria and tetanus	age of 11yr-up to 13yr	once: age of 11yr	at the age of 11yr (1 coupon)
MR measles-rubella	primary: age of 1yr-up to 2yr ----- booster: the year before entering school (Apr.-Mar.)	once: 12mo-up to 24mo age of 1yr ----- once: the year before entering school (Apr.-Mar.)	at the age of 1yr (1 coupon) ----- April of 1 yr before entering school (1)
Varicella	age of 12mo-36mo	twice: age of 12-15mo/ after 6mo-12mo of interval	at the age of 1yr (2 coupons)
Japanese Encephalitis: primary period	initial: age of 6mo-up to 90mo ----- booster: age of 3mo-up to 90mo	twice: age of 3yr (1-4wk interval each) ----- once: age of 4yr/ 1yr after completing initial inoculation	at the age of 3yr (2 coupons) ----- at the age of 4yr (1 coupon)
secondary period	age of 9yr-up to 13yr	once: age of 9yr	inform Health Center if desired
(special case)	people who were born between April 2, 1995 and April 1, 2007 can receive it until they reach the age of 20yr as a special aid recipient		
HPV (bivalence or quadrivalence selectable) human papillomavirus vaccine	girls about age of 12-16yr (girl students between the 6th grade at primary school and the 1st grade at high school)	bivalence: 3 times: 0, 1mo and 6mo ----- quadrivalence: 3 times: 0, 2mo and 6mo	inform Health Center if desired