

GOVERNMENT SUBSIDIZED INOCULATION

as of March 2020

vaccine	aid recipients	dose	standard inoculation age	Qpon delivered
HIB hemophilus influenza type b vaccine	2mo-up to 5yr	initial x 3	age of 2mo-6mo (4-8week-interval each)	shortly before the age of 2mo
		booster x 1	7-13mo after completing initial inoculation	
PCV-13 pneumococcal polysaccharide conjugated vaccine	2mo-up to 5yr	initial x 3	age of 2mo-6mo (4week-interval each)	shortly before the age of 2mo
		booster x 1	age of 12mo-15mo 6mo after completing initial inoculation	
HEPATITIS B	up to 1yr	initial x 2 last	age of 2mo-9mo (4wk-interval) 20weeks after the 1st shot	shortly before the age of 2mo
BCG (tuberculosis) bacille de calmette guerin	up to 1 yr	1 dose	age of 5-8mo	shortly before the age of 3mo
DPT-IPV : primary period (diphtheria, tetanus and acellular pertussis-IPV)	age of 3mo-90mo	initial x 3	age of 3mo-12mo (3-8wk-interval each)	shortly before the age of 3mo
		booster x 1	6mo, usually 12-18mo after completing initial inoculation	shortly before the age of 18mo
IPV (inactivated polio virus) DPT-IPV for recipient born after August 2012	age of 3mo-90mo	initial x 3	age of 3mo-12mo (3wk-interval each)	inform Local Health Center, if desired
		booster x 1	6mo, usually 12-18mo after completing initial inoculation	
MR (measles-rubella) primary period	age of 12mo-24mo	1 dose	age of 12mo (soon after age of 12mo)	shortly before the age of 1yr
	secondary period	the year before entering primary school	1 dose	the year before entering primary (April-March)
VARICELLA	age of 12mo-36mo	1st dose	age of 12mo-15mo	shortly before the age of 1yr
		2nd dose	3mo, usually 6-12mo after 1st dose	
JAPANESE ENCEPHALITIS primary period	age of 6mo-90mo	initial x 2	at age of 3yr (1-4wk-interval)	shortly before the age of 3yr
		booster x 1	at age of 4yr 6mo, usually 1yr after completing initial inoculation	shortly before the age of 4yr
secondary period	age of 9yr-13yr	1 dose	at age of 9yr	shortly before 9yr
special case	recipients born between April 2, 1999 and April 1, 2007 can receive all the doses until they reach the age of 20yr			
	recipients born between April 2, 2007 and Oct. 1, 2009 can receive the dose of primary period (3 doses) at the age of 9yr-13yr			
DT : secondary period (diphtheria and tetanus)	age of 11yr-13yr	1 dose	age of 11yr	shortly before the age of 11yr
HPV (human papiloma virus) bivalence or quadrivalence selectable	girls about age of 12-16yr	bivalence 3 dose	0, 1mo and 6mo	inform Local Health Center, if desired
		quadrivalence 3 dose	0, 2mo and 6mo	