

QUADRIVALENT VACCINES FOR USE IN THE 2020-2021 INFLUENZA SEASON

A/ Guangdong-Maonan/ SWL 1536/2019 (H1N1) pdm0 B/ Phuket/ 3073/ 2013 (Yamagata lineage)
A/ Hong Kong/ 2671/2019 (H3N2) B/ Victoria/ 705/ 2018

RECIPIENT TO COMPLETE

First shot Second flu shot for this season

Family name _____ First name _____ DOB _____

PLEASE ANSWER YES OR NO FOR EACH QUESTION

		Remarks
1. Do you have any severe allergies to anything, including eggs or latex?	<input type="checkbox"/> No <input type="checkbox"/> Yes	
2. Do you have a history of Guillan-Barre Syndrome, a severe paralyzing illness?	<input type="checkbox"/> No <input type="checkbox"/> Yes	
3. Have you been vaccinated against flu in previous years?	<input type="checkbox"/> No <input type="checkbox"/> Yes	
4. Do you suffer from any medical conditions, or are you taking any medications?	<input type="checkbox"/> No <input type="checkbox"/> Yes	
5. Have you ever had a severe reaction following any vaccine?	<input type="checkbox"/> No <input type="checkbox"/> Yes	
6. Do you feel unwell today?	<input type="checkbox"/> No <input type="checkbox"/> Yes	
7. (Women only) Are you pregnant?	<input type="checkbox"/> No <input type="checkbox"/> Yes	

GENERAL INFORMATION ABOUT INFLUENZA

The flu virus infects the nose, throat and lungs and can cause various complications, most commonly ear infections and pneumonia.

It is spread through the air by coughing and sneezing and the disease is caught by inhaling the airborne virus. Flu symptoms include the abrupt onset of fever (lasting 3-4 days), headaches, fatigue (tiredness), muscle or body aches, sore throat, runny or stuffy nose, cough.

The best way to prevent seasonal flu is to get vaccinated every year. Flu vaccination can help protect yourself from flu and also helps protect the people around you who are more vulnerable to flu and its complications.

It takes about 2 weeks after vaccination for antibodies that protect against flu to develop in the body, and lasts for 4-5 months.

CAUTIONS AFTER RECEIVING THE VACCINATION

Minor problems following flu shot include: soreness, redness or swelling at the injection site, low-grade fever, runny nose, nasal congestion, cough, headache, tiredness.

(If they occur, usually starts after the shot and lasts 1-2days)

Can take a pain killer, such as paracetamol or ibuprofen to ease the discomfort, but DO NOT TAKE ASPIRIN.

Safe to carry out normal daily activities, but avoid strenuous exercise and excessive alcohol intake.

Can take a bath or shower, but do not scrub the injection site.

CONSENT

By signing this form I agree that I have read and understood the information above.

I give my consent to receive the influenza vaccination today.

Signature of recipient(Guardican, if applicable) _____ Date _____

MEDICAL STAFF USE ONLY

Route: SC

Dose: 0.5ml 0.25ml (I / I · I / II · II / II)

Site: Upper Arm Right Left

Thigh / Buttock Right Left

Manufacturer Daiichi Sankyo KMB

Handai Biken Denka

Batch Number: Kitasato

Signature of vac. Giver _____